

SYNTHESIS OF 1-(6-CHLORO-2-METHYL-4-PHENYLQUINOLINE) ETHANONE USING DIFFERENT HETEROGENEOUS CATALYSTS IN DRY MEDIA UNDER MICROWAVE IR-RADIATION

Pooja

poojamaharia91@gmail.com
CSIR NET, Chemistry

Received: May 23, 2018

Accepted: July 02, 2018


ABSTRACT

1-(6-Chloro-2-methyl-4-phenylquinoline) ethanone is synthesized in dry media using heterogeneous catalyst in high yield in shorter reaction time under microwave ir-radiations.

Keywords: Indole, Microwave, Aryl, heterocyclic.

Introduction:

The trisubstituted quinolines have wide range of biological activities as anti-malarial, anti-bacterial, anti-asthmatic, anti-hypertensive, anti-inflammatory, anti-platelet activity and as tyro-kinase PDGF-RTK inhibiting activity.¹⁻³ The synthesis of trisubstituted quinolines under conventional refluxing conditions require longer reaction time and tedious work up so, there existed a need for alternative methods to carry out the synthesis of trisubstituted quinolines. Microwave assisted reactions are gaining much more importance in synthetic organic chemistry due to dramatic reduction in time from days to hours and hours to minutes or seconds.⁴⁻²⁴ The conventional heating reaction conditions are modified by changing media and catalyst. The present work reports the synthesis of 1-(6-chloro-2-methyl-4-phenylquinoline) ethanone in dry media using heterogeneous catalyst in high yield in shorter reaction time under microwave irradiations (Scheme-1).



(Scheme)

We initiated our investigations by condensing acetyl acetone (1mmole) with 5-chloro-2-aminobenzophenone (1mmole) at 80 W, 160 W, 240 W, 320 W, 400 W, 480 W and 560 W in the presence of Mont.K-10. The results obtained are shown in Table-1 below. As can be seen from the Table-1 that when ethylacetoacetate (1mmole) react with 5-chloro-2-aminobenzophenone (1mmole) to give 1-(6-Chloro-2-methyl-4-phenylquinoline) ethanone, 560 W power level proved to be the best from the yield point of view.

Table-1: Synthesis of 1-(6-Chloro-2-methyl-4-phenylquinoline) ethanone using Mont.K-10 under various irradiation (power levels).

Sr. No.	Power Levels (watts).	Yield (%).	Time (mints.)
1	80	83	5
2	160	85	5
3	240	87	5
4	320	89	5
5	400	92	5
6	480	94	5
7	560	96	5

We next carried out the formation of 1-(6-Chloro-2-methyl-4-phenylquinoline) ethanone by condensing acetylacetone (1mmole) with 5-chloro-2-aminobenzophenone(1mmole) at 560W in the presence of Mont. KSF, silica gel, MgSO₄ (anhyd), Na₂SO₄ (anhyd), Yb(OTf)₃, Sc(OTf)₃, Dy(OTf)₃, Gd(OTf)₃, InCl₃, Y(OTf)₃ and Bi(OTf)₃ catalysts. The results obtained have also been collected in the Table 2.

Table-2: Synthesis of 1-(6-Chloro-2-methyl-4-phenylquinoline) ethanone using different catalysts at 560W under various ir-radiation.

Sr. No.	Catalyst	Time (in min.)	Yield (%)
1	Mont. K-10	5	96
2	Mont. KSF	5	95
3	Silica gel	5	94
4	MgSO ₄ (anhyd)	5	93
5	Na ₂ SO ₄ (anhyd)	5	92
6	Yb(OTf) ₃	5	91
7	Sc(OTf) ₃	5	90
8	Dy(OTf) ₃	5	89
9	Gd(OTf) ₃	5	88
10	InCl ₃	5	87
11	Y(OTf) ₃	5	86
12	Bi(OTf) ₃	5	85

Experimental:

All the melting points reported are uncorrected. Infrared spectra (V_{max} in cm⁻¹) were recorded in nujol mull or KBr on a Perkin-Elmer 842/Beckman IR-20 / Hitachi 215 spectrometers. The proton magnetic resonance spectra were recorded on a VXR-200 MHz or R-32 Perkin-Elmer 90 MHz spectrometer in CDCl₃ or DMSO-d₆ using tetramethylsilane (TMS) as internal reference standard. The chemical shifts are expressed in δ (ppm) units downfield from TMS. Mass spectra were scanned on a Jeol JMX-DX-300 spectrometer operating at 70 eV. Carbon, hydrogen and nitrogen analyses were carried out on a Yanaco MT -3 (JAPAN) instrument. Thin layer chromatography (TLC) were performed on silica-gel plates using acetone-benzene (1 : 3 or 1 : 2) as solvent system and iodine chamber as visualizing agent.

Typical procedure for the synthesis of 1-(6-Chloro-2-methyl-4-phenylquinoline) ethanone: A mixture of acetylacetone (1mmole), 5-chloro-2-aminobenzophenone (1mmole), catalyst (1g) was taken in an Erlenmeyer flask (100 ml) and was ir-radiated for 5 minutes at 70% power level (560 W) in an unmodified domestic microwave oven operating at 2450 MHz. After cooling to room temperature, the crude product was extracted and recrystallised with ethanol to yield pure 1-(6-Chloro-2-methyl-4-phenylquinoline) ethanone. Mp observed : 150°C, reported: 150°C : 1H - NMR (δ in ppm in CDCl₃): δ =2.0 (s, 3H), 2.7 (s, 3H), 7.25-7.40 (m, 2H), 7.5-7.6 (m, 3H), 7.7 (m, 2H), 8.0 (d, J = 8 Hz, 1H) and IR (KBr, cm⁻¹): 3026, 2950, 1695, 1605, 692.

References:

- Larsen, R. D.; Corley, E. G.; King, A. O.; Carrol, J. D.; Davis, P.; Verhoeven, T. R.; Reider, P. J.; Lablbe, M.; Gauthier, J. Y.; Xiang, Y. B.; Zamboni, R. J. *J. Org. Chem.* 1996, 61, 3398. (b) Chen, Y. L.; Fang, K. C.; Sheu, J. Y.; Hsu, S. L.; Tzeng, C. C. *J. Med. Chem.* 2001, 44, 2374. (c) Roma, G.; Braccio, M. D.; Gmattioli, G. F.; Ghia, M. *Eur. J. Med. Chem.* 2000, 35, 1021.
- Kalluraya, B.; Serrnivas, S.; Farmaco 1998, 53, 399. (b) Doube, D.; Blouin, M.; Brideau, C.; Chan, C.; Desmarais, S.; Eithier, D.; Fagueyret, J. P.; Friesen, R. W.; Girard, M.; Girard, Y.; Guay, J.; Tagari, P.; Young, R. N., *Bioorg. Med. Chem. Lett.* 1998, 8, 1255.
- Ko, T. C.; Hour, M. J.; Liem, J. C.; Teng, C. -M.; Lee, K -H.; Kuo, S.-C.; Huang, L.-J.; *Bioorg. Med. Chem. Lett.* 2001, 11, 279. (b) Ferrarini, P. L.; Mori, C.; Badwneh, M.; Manera, C.; Martinelli, A.; Miceli, M.; Ramagnoli, F.; Saccomanni, G. *J. Hetetocycl. Chem.* 1997, 34, 1501. (c) Maguire, M. P.; Sheets, K. R.; Mcverty, K.; Spada, A. P.; Zilberstain, A. J. *Med. Chem.* 1994, 37, 2129.
- Anastas, P.T. and Farris, C.A. (Eds.), *Benign by Design: Alternative Synthetic Design for Pollution Prevention*, ACS symposium, Ser. N. 557. Washington DC, 1994.
- Collins, T. *Towards sustainable chemistry*, Science, 2001, 291, 5501, 48.
- Anastas, P. and Warner, J.C. *Green Chemistry: Theory and Practice*, Oxford Science Publicatons, Oxford, 1998.
- Collins, T.J. *Green Chemistry*, Macmillan, Encyclopedia of Chemistry, New York, 1997.