

Surfactants as penetration enhancer in transdermal drug delivery system

Churala Pal¹ & Prabal Giri^{2*}

¹Department of Chemistry, Basanti Devi College, Kolkata 700029,
West Bengal, INDIA.

^{2*}Department of Chemistry, Gushkara Mahavidyalaya, Burdwan 713128,
West Bengal, INDIA.

Received June 22, 2017

Accepted July 19, 2017

ABSTRACT

Human skin is a remarkably efficient barrier, designed to keep “our insides in and the outsides out”. This barrier property causes difficulties for transdermal delivery of therapeutic agents. One long-standing approach to increase the range of drugs that can be effectively delivered via this route has been to use penetration enhancers, chemicals that interact with skin constituents to promote drug flux.

To-date, a vast array of chemicals has been evaluated as penetration enhancers (or absorption promoters), yet their inclusion into topical or transdermal formulations is limited since the underlying mechanisms of action of these agents are seldom clearly defined. In this article we review some uses of the more widely investigated chemical penetration enhancers and discuss possible mechanisms of action.

Key Words: Transdermal, Surfactant, Penetration enhancer, Skin,

Introduction

Transdermal drug delivery is the topical application of drugs to the skin in the treatment of skin diseases, wherein high concentrations of drugs can be localized at the site of action, thereby reducing the systemic drug levels and side effects¹⁻³. 'U.S. Emerging Transdermal Drug Delivery Technologies Markets', reveals that this market generated revenues worth \$1.57 billion in 2002 and reached a staggering \$5.67 billion in 2009⁴. In 1924, Rein proposed that a layer of cells joining the STRATUM CORNEUM — the thin, outermost layer of the skin — to the EPIDERMIS posed the major resistance to transdermal transport⁵. The corneocytes, which comprise cross linked keratin fibres, are about 0.2–0.4 μm thick and about 40 μm wide⁶. Penetration enhancers are used to promote the drug transport across the skin barrier. The interaction of the enhancers with the polar head groups of the lipids is the possible way to increase the penetration⁷. Surfactants have the potential for the solubilization of the stratum corneum lipids and thus act as penetration enhancers. Keratin interactions are also thought to explain the penetration-enhancing effects of surfactants⁸.

Target site for Transdermal drug delivery system - Skin

The outermost layer of the epidermis, the stratum corneum, provides a formidable barrier to dermal absorption that determines the rate of dermal penetration⁹⁻¹⁴. The stratum corneum differs from the rest of the epidermis in being a two-compartment tissue consisting of dead cornified cells (corneocytes) with a matrix of intercellular lipids¹⁵. The hydrophilic properties of the skin increase as the depth increases from the surface, such that the viable epidermis represented by the stratum granulosum, the stratum spinosum and the stratum basale, respectively is significantly hydrophilic. The dermis layer is also hydrophilic, hence favoring the uptake of hydrophilic chemicals¹⁶. The viable epidermis contains corneocytes at varying stages of differentiation, as well as melanocytes, Langerhans cells (important for antigen presentation and immune response), and Merkel cells (involved in sensory perception). This layer facilitates the diffusion of, for example, xenobiotics and decreases in surface area with age¹⁷.




Fig (1): Diagrammatic representation of skin

Barriers posed by skin against percutaneous absorption

Corneocytes are the 'bricks' embedded in an intercellular lipid matrix of mainly fatty acids, ceramides, cholesterol and cholesterol sulfate¹⁸. The corneocytes are held together by corneodesmosomes, which confer structural stability to the stratum corneum. The stratum corneum lipids are composed primarily of ceramides, cholesterol and fatty acids that are assembled into multi-lamellar bilayers. This unusual extracellular matrix of lipid bilayers serves the primary barrier function of the stratum corneum¹⁹. The cells are joined together by desmosomes, maintaining the cohesiveness of this layer²⁰. The heterogeneous structure of the stratum corneum is composed of approximately 75-80% protein, 5-15% lipid and 5-10% unidentified on a dry weight basis²¹. There are two general options for drug substances to permeate the stratum corneum: the transepidermal route and the route via pores²².

Factors affecting skin penetration:

- Thickness of horny layer
- Skin condition

Factors associated with medicament:

- Solubility
- Dissociation constant
- Particle size

Factors associated with vehicle:

- Contact with skin
- Penetration into epidermis
- Alteration of skin permeability

ROUTES OF DRUG PERMEATION THROUGH THE SKIN.

1. Intercellular route
2. Transcellular route
3. Follicular route

1. Intercellular route: The more common pathway through the skin is via the intercellular route.

Drugs crossing the skin by this route must pass through the small spaces between the cells of the skin.

2. Transcellular route: Drug crossing the skin via this route must pass through the cells. (keratinocytes)

3. Transappendagal route: Passage of molecules via sweat glands, hair follicles and sebaceous glands.

Penetration Enhancers

Currently, the most widely used approach to drug permeation-enhancement across the stratum corneum barrier is the use of chemical penetration enhancers (sorption promoters and accelerants). One of the most recent comprehensive reviews on the classes of enhancers used was written by Ghosh et al²³. According to Shah, enhancers:

- increase the diffusivity of the drug in the skin;
- cause stratum corneum lipid-fluidization, which leads to decreased barrier function (a reversible action);
- increase and optimize the thermodynamic activity of the drug in the vehicle and the skin;
- result in a reservoir of drug within the skin;
- affect the partition coefficient of the drug, increasing its release from the formulation into the upper layers of the skin²⁴.
- disrupt the order within and between the corneocyte upon binding to the keratin filament²⁵

The following classes of compounds have been tested for their enhancer action: water, hydrocarbons (alkanes and alkenes); alkanols and alkenols; acids; esters; alkyl amino esters; amides; ureas; amines and bases; sulfoxides; terpenes²⁶, steroids; dioxolanes; pyrrolidone and imidazole derivatives; laurocapram (Azone) and its derivatives. Other approaches to enhancement include the use of enzymes, natural oils, phospholipid micelles, liposomes, niosomes, polymers²⁷⁻³⁰, isopropyl myristate³¹, nicotinic acid esters³², ethanol, hydrogenated Soya phospholipid³³, essential oils³⁴, n-octanol and decanol³⁵, surfactants³⁶⁻⁴³ have been reported to enhance the permeability of drugs.

Surfactants

Surfactants are frequently used as emulsifiers in formulations for dermal application. A substance which is positively adsorbed at the liquid/vapour and/or at other interfaces are called surfactants⁴⁴. Surfactants are usually organic compounds that are amphiphilic, meaning they contain both hydrophobic groups (their tails) and hydrophilic groups (their heads). Therefore, a surfactant molecule contains both a water insoluble (or oil

soluble) component and a water soluble component. Surfactant molecules will diffuse in water and adsorb at interfaces between air and water or at the interface between oil and water, in the case where water is mixed with oil⁴⁵.

Classification of surfactants

Surfactants can be classified into four main categories according to the presence of formally charged groups in the head; anionic (e.g. sodium laurylsulfate), cationic (e.g. cetyltrimethyl ammonium bromide), nonionic (e.g. polyoxyethylene sorbitan monopalmitate) and amphoteric (e.g. N-dodecyl-N, N-dimethylbetaine). The investigation of enhancing abilities of nonionic surfactants has been focused on five principal series of surfactants, which are polysorbates, sorbitan esters, polyoxyethylene alkylethers, polyoxyethylene alkylphenols and poloxamers⁴⁶. It is generally recognized that nonionic surfactants possess the least toxicity and skin irritation potential⁴⁷, and therefore they have been widely investigated as skin penetration enhancers.




Fig (3): Classification of Surfactants

Mechanism of action of surfactants as penetration enhancers

Anionic Surfactants

In general, anionic surfactants are more effective than cationic and nonionic surfactants in enhancing skin penetration of target molecules. Some anionic surfactants interact strongly with both keratin and lipids. alter the permeability of the skin by acting on the helical filaments of the stratum corneum, thereby resulting in the uncoiling and extension of keratin filaments to produce keratin. Then they cause an expansion of the membrane, which increases permeability⁴⁸.

Sodium lauryl sulfate (SLS), an anionic surfactant, possesses skin penetration enhancer properties and enhances penetration into the skin by increasing the fluidity of epidermal lipids⁴⁹⁻⁵². An additional mechanism for the skin penetration enhancement by SLS could involve the hydrophobic interaction of the SLS alkyl chain with the skin structure which leaves the end sulphate group of the surfactant exposed, creating additional sites in the membrane which leads to permit an increase in skin hydration^{53,54}.

Cationic Surfactants

The cationic surfactants interact with the keratin fibrils of the cornified cells and result in a disrupted cell-lipid matrix. The cationic surfactants may interact with anionic components of the stratum corneum, change the electronic property there, and stimulate the transfer of the anionic drug into the skin⁵⁵.

Nonionic Surfactants

The nonionic surfactants enhance absorption by inducing fluidization of the stratum corneum lipids⁴⁸.

They are two possible mechanisms by which the rate of transport is enhanced using nonionic surfactants.

Initially, the surfactants may penetrate into the intercellular regions of SC, increase fluidity and eventually solubilise and extract lipid components.

Secondly, penetration of the surfactant into the intercellular matrix followed by interaction and binding with keratin filaments may result in a disruption within the corneocyte^{47, 56}.

Factors governing the activity of Surfactant as penetration enhancer

Critical micelle concentration

In biological systems the effects of surfactants are complex, particularly their effect on cell membranes, which can lead to alterations in permeability⁴⁶. The effect of surfactants on

membrane permeability describe an apparent concentration-dependent biphasic action, such that an increase in membrane permeability occurs at low surfactant concentrations, but this decreases at higher concentrations, generally above the critical micelle concentration (CMC) of the surfactant. Above the CMC, the added surfactant exists as micelles in the solution and micelles are too large to penetrate the skin⁴⁸. The CMC represents a narrow range of concentrations above which surfactants form dynamic aggregates known as micelles. The structure of micelles is such that in aqueous solution the monomers are aligned with their hydrophobic regions towards the centre and their hydrophilic sections outwards towards the aqueous bulk⁵⁷.

Chain length of carbon atoms

Enhancement depends on the carbon chain structure of the enhancer⁵⁸⁻⁶⁰. Maximal enhancement is generally attained for enhancers with a carbon chain length in the range of 10–14. This optimal range was found for anionic, cationic and neutral enhancers⁶¹⁻⁶⁵.

Transdermal gradient

The driving force for penetration into the skin is the “ transdermal gradient” caused by the difference in water content between the relatively dehydrated skin surface (approx 20% water) and the aqueous viable epidermis (close to 100%)⁶⁶.

Hydrophilicity of surfactant head (Laughlin's hypothesis)

Surfactants with hydrophilic head groups should more effectively enhance the percutaneous penetration of polar molecules, while those of lesser hydrophilicity should be less effective The results obtained in the present work in agreement with Laughlin's hypothesis because Cetyltrimethylammonium bromide ($\log P_{oct} < 1$) which is more hydrophilic than benzalkonium chloride ($\log P_{oct} = 1.9$) is less effective in enhancing lorazepam skin penetration. This could be attributed to the lipophilicity of lorazepam⁶⁷.

Steric forces

Steric repulsive forces are caused by the reduced conformational freedom of adsorbed molecules and changes in molecule/solvent interactions as two surfaces are approached. They are present in both surfactant and polymer systems and increases in magnitude and range with the size of the adsorbed molecules⁶⁸.

PERMEANT	SURFACTANT	DESCRIPTION	REFERENCES
Chloramphenicol	Sodium Lauryl Sulphate	significant enhancement of surfactant facilitated permeation through hairless mouse skin	36,37
Hydrocortisone, lidocaine	Tween 80	Acceleration of permeating across hairless mouse skin	36,37
5-fluorouracil, antipyrine and 2-phenyl ethanol	Span 20 (1 and 5% w/v in ethanolic solution)	increased the penetration through Wistar rat epidermis in- vitro	40
Naproxen, naloxone	sodium decyl and dodecyl sulfates	increased the in vitro permeation rates	69,61
Lorazepam	Sodium lauryl sulfate 5%, Cetyltrimethylammonium bromide	enhancing activity on the skin permeation across rat skin in vitro	70
Lidocaine (from saturated systems in propylene glycol-water mixtures)	Cationic surfactants	promote the permeation through excised human skin	71
5-flourouracil	0.1% Tween 20 in normal saline	permeation across hairless mouse skin 6-fold	72
Chloramphenicol	0.5 and 1% Tween 80	increased the skin penetration	73
Laurocapram	SLS	produce variations in the structural organisation of lipids when it is used above the critical micellar oncentration ⁷³	75,76
Fluoxetine	Labrasol (caprylo-capryloyl macrogol-8-glyceride)	Permeation significantly enhanced from a vehicle system consisting of 65% v/v ethanol	77
Nitrendipine	Benzalkonium chloride, SLS, Tween 80	enhance the permeation of nitrendipine across rat skin.	78
Piroxicam	polyoxyethylene-2-oleyl ether	enhancing effects with an enhancement factor of 2.84.	79
Diazepam	sodium lauryl sulfate	increase the permeation of diazepam across rat skin	43
Foscarnet	sodium lauryl sulfate (SLS)	increase the penetration thereby, its efficacy against HSV-1 cutaneous lesions in hairless mice.	80
Hydrocortisone	Span 20: Span 80	an increase in diffusion through the skin (58.29%)	81
Mepivacaine	polyoxyethylene 2-oleyl ether	enhancement of permeation through skin	82
Diltiazem hydrochloride	Tween80	improve the in vitro permeation through pig ear skin	83
Hydrocortisone Acetate	2-(2-ethoxy-ethoxy) ethanol (Trans-Cutol)	Functions as solubilizer and permeation enhancer.	84
Acyclovir	Transcutol	enhancing ability to pass through the skin	85,86,87
Progesterone	medium-chain mono- and diglycerides	ability to interact with membrane lipids and proteins, increasing membrane permeability enhanced the topical and transdermal delivery of PGT by 2.5- and 7-fold	88,89,90,91
Adenosine	polysorbate 80, medium-	increases in the skin	90,92,93,94

	chain glycerides, and propylene glycol	penetration and transdermal delivery	
Fluconazole	Labrasol (Lab) /EtOH (1:1, w/w) mixture	showed the highest permeation profile among other formulations 9.12 times higher	95,96
Ropinirole	Tween 20	the skin permeation increased from 20% to 35%	97,98
Flurbiprofen	span 20, span 80	showed a larger amount of flurbiprofen penetration through rabbit skin	99
5-fluorouracil	lecithin/ethanol/decyl glucoside (14.67:12.15:18.18% w/w.)	significant increase in permeability coefficient in newly born mice skin	100
Interferon	nonoxynol-9	indications such as ending of new-lesion formation, scabbing, and healing of lesions in patients with recurrent genital herpes	101
Piroxicam	Tween 80:Span 20	enhance permeation through various skin models by reversible disturbance of the stratum corneum layer	40,79,102,103
Paclitaxel	Cremophor EL	intracellular penetration enhancer for liposomal drugs to improve their anticancer efficiency	104
Ibuprofen	Tween 80	Higher penetration through the skin	105
2,3,5,6-tetramethylpyrazine	Labrasol®	the skin penetration were 0.9- to 4.7-fold higher	106
Ketorolac	Span 60, Tween 20	provided a higher ketorolac flux across the skin (7-and 4-fold the control, respectively)	107
Amlodipine besilate	SLS, tween 20, sodium tauroglycolate and hyaluronidase	hyaluronidase enzyme show higher permeability and steady state flux increased linearly with increasing donor concentration	108,109
Lacidipine	Tween 80 and Labrasol	enhances the drug permeation through the skin	110
Nicardipine Hydrochloride	Tween 80 / Span 20 mixture	enhances the drug permeation through the skin	111
Ramipril	Cremophor EL	enhances the drug permeation through the skin	112
Tamoxifen citrate	Cremophore EL, Tween 80	excellent skin penetration rate of drug	113
Diclofenac diethylamine	Span 20	enhancement of skin permeation of the drug of up to 30% in rat skin	114
Atenolol	Polyoxyethylene-2-oleyl ether	best enhancement for the atenolol transdermal drug delivery	115
Propranolol, metaproterenol sulfate	Lauric acid	enhancement on the percutaneous absorption	116, 117
Methyl nicotinate	polyoxyethylene (10) lauryl ether, polyoxyethylene (10) stearyl ether	permeability rate of methyl nicotinate from aqueous solution is increased two-fold	118

Various surfactants used to enhance penetration across the skin in current scenario

Tween 80

Acceleration of hydrocortisone and lidocaine permeating across hairless mouse skin by the nonionic surfactant Tween 80^{36,37}. It has been shown that at concentrations of 0.5 and 1% Tween 80 increased the skin penetration of chloramphenicol⁷³. It is apparent that propylene glycol and Tween 80 interact to affect the skin barrier so as to promote the penetration of lorazepam. It was evident from surface tension studies that the addition of propylene glycol raises the CMC of the nonionic surfactants by approximately a factor of 10. The increase in monomer concentration might be an explanation for observed synergistic effect of propylene glycol and Tween 80. Highest permeation rate was observed with the solution containing 1% w/w of Tween 80 in diazepam permeation⁴³. Initially, the surfactants may penetrate into the intercellular regions of stratum corneum, increase fluidity and eventually solubilise and extract lipid components. Secondly, penetration of the surfactant into the intercellular matrix followed by interaction and binding with keratin filaments may results in a disruption within the corneocyte. Tween 80 is thought to enhance the penetration of lorazepam via both the lipophilic and the hydrophilic molecular mechanisms, and to disrupt the lipid arrangements in the stratum corneum and to increase the water content of the proteins in the barrier^{56,57}. The structure of Tween 80 is relevant to this role. It contains the ethylene oxide and a long hydrocarbon chain. This structure imparts both lipophilic and hydrophilic characteristics to the enhancer, allowing it to partition between lipophilic mortar substance and the hydrophilic protein domains. Tween 80 may interact with the polar head groups of the lipids and the modification of H-bonding and ionic forces may occur. The other possible mechanism related to our studies involves the protein domains (keratinocytes). In this case, targets of the enhancer are the keratin fibrils and their associated water molecules. The disruption caused by the enhancer makes this area more aqueous. With high enough volumes an increase in the solubilising ability of the aqueous layer could result and actually change the operational partition coefficient of this region of the skin⁷². This would then allow for drug transport through the corneocytes.

Sodium Lauryl Sulfate

Surfactant facilitated permeation of many materials through skin membranes has been researched, with reports of significant enhancement of materials such as chloramphenicol through hairless mouse skin by sodium lauryl sulfate,^{36,37}. Sodium lauryl sulfate at 5% showed a remarkable enhancing activity on the skin permeation of lorazepam across rat skin in vitro. A marked increase in the drug flux was attributed to the skin damage caused by this anionic surfactant at 5% concentration, the highest concentration used in the study⁷⁰. Sodium lauryl sulfate is able to produce variations in the structural organisation of lipids when it is used above the critical micellar concentration⁷⁴, and similar effects on organisation of skin lipids have been described for other permeation enhancers such as Laurocapram^{75,76} reported that SLS was able to increase the penetration rates of compounds that have values of lipophilicity lower than an optimum lipophilicity. An additional mechanism for the skin penetration enhancement by SLS could involve the hydrophobic interaction of the SLS alkyl chain with the skin structure which leaves the end sulphate group of the surfactant exposed, creating additional sites in the membrane which leads to permit an increase in skin hydration^{53,54}.

Dodecyltrimethylammonium bromide

Dodecyltrimethylammonium bromide (DTAB) as to the pretreatment with cationic surfactant DTAB, opposing effects on the flux are found compared to LA. During all three experimental intervals (passive before iontophoresis, iontophoresis, passive after iontophoresis) an inhibition. This is most likely related to the positive charge of surfactant DTAB. During the passive period, the partitioning of the positively charged R-apomorphine into the membrane is hindered by the repulsion of absorbed positively charged DTAB. After turning on the current, DTAB is driven into the skin and compensates for the native negative charge of human stratum corneum, thereby reducing the electro-osmotic flow¹¹⁹

Laureth-3 oxyethylene ether

Laureth-3 oxyethylene ether (C₁₂EO₃) the nonionic surfactant C₁₂EO₃ substantially increased iontophoretic transport rate of R-apomorphine by 2.3-fold, whereas passive delivery was basically unchanged or slightly affected. The magnitude of enhancing effect of C EO was dependent on the

surfactant concentration and the pretreatment duration¹²⁰.

Span 20

Pretreatment of skin with Span 20 (1 and 5% w/v in ethanolic solution) significantly increased the penetration of 5-fluorouracil, antipyrine and 2-phenyl ethanol through Wistar rat epidermis in vitro⁴⁰.

Tween 20

Tween 20 has been shown to increase the permeation of hydrocortisone and lidocaine across hairless mouse skin in vitro^{36,37}.

Sodium Lauroyl sarcosinate & Sorbitan Monolaurate

Sodium lauroylsarcosinate, and a nonionic surfactant, sorbitan monolaurate, more markedly increased the transdermal flux of drugs than the individual components used alone. Moreover, the formulation exhibited a reduction in skin irritation¹²¹.

Sodium Decyl & Dodecyl Sulfates

Sodium decyl and dodecyl sulfates increased the in vitro permeation rates of several drugs including naproxen¹²² and naloxone^{48,60} reported that the capacity of the stratum corneum to retain significant quantities of membrane-bound water is reduced in the presence of sodium dodecanoate and sodium dodecyl sulfate. This effect is readily reversible upon removal of the agents. These investigations proposed that anionic surfactants alter the permeability of the skin by acting on the helical filaments of the stratum corneum, thereby resulting in the uncoiling and extension of β -keratin filaments to produce α -keratin. Then they cause an expansion of the membrane, which increases permeability.

Cetyltrimethylammonium bromide

The permeation profile of lorazepam in presence of the other cationic surfactant, CTAB, reveals that an increase in the concentration of CTAB cetyltrimethylammonium bromide results in an increase in the flux of lorazepam. Similar results were reported on the effect of other cationic surfactant cetrimide which is a cationic surfactant which contains higher percentages of CTAB on haloperidol permeation through rat skin¹²³.

n-Dimethyldialkylammoniums

Enhancement effects of the double-chained cationic surfactants of n-dimethyldialky

lammoniums $(CH_3)_2N_1(C_nH_{2n+1})_2$ on the permeation of anionic salicylate through excised guinea pig dorsal skin at pH 7.4. n-dimethyldidecylammonium (2C10), which seemed to form micelles, had dose-dependent enhancement effects and about a ninety-fold increase in the permeability. Dimethyldilaurylammonium (2C12), seemed to form bilayer vesicles, induced about a twenty five-fold increase in the permeability^{124, 125, 126}.

Sodium Dodecyl Sulfate

Sodium dodecyl sulfate (SDS) and dodecyl trimethylammonium bromide (C12TAB) Patist et al. have previously shown that SDS micellar stability may be tailored by the addition of oppositely charged surfactants such as alkyltrimethylammonium bromides (C_n TABs). The long-chain TABs enhance SDS micellar stability, as measured by relaxation time, by up to 2000 times. addition of C_{12} TAB to SDS leads to stabilization of micelles and sub-micellar aggregates and such stabilization decreases and even virtually eliminates sub-micellar aggregates^{127, 128}.

Polyoxyethylene-23-lauryl ether, polyoxyethylene-2-oleyl ether and polyoxyethylene-2-stearyl ether

Poloxamer gels containing piroxicam including surfactants as enhancers are good preparations to promote the percutaneous absorption of drugs¹¹⁵.

Cremophor RH 40®

Cremophor RH 40® shifts the drug distribution to the stratum corneum. Cremophor RH 40® enhanced the flufenamic acid content in the stratum corneum 2-fold. The amount of flufenamic acid after pretreatment with Cremophor RH 40® is on a higher level in the stratum corneum¹²⁹.

Propylene glycol, 2-(2-ethoxy-ethoxy) ethanol (Transcutol®)

The optimum formulation containing 2.5% Transcutol as the penetration enhancer shows 1.7-fold enhancement in flux and permeation coefficient as compared to marketed cream and ointment formulation. In order to further improve the permeation rate of Acyclovir from the microemulsion, enhancer like Transcutol in the concentration ranging from 1% to 5% was employed. Results indicate significant improvement in the permeation pattern of ACV with the incorporation of enhancers. The presence of Transcutol in the formulations also results in an increase in the mean cumulative amount. The

enhancing ability of Transcutol has been attributed to its ability to pass through the skin and get incorporated into the multiple-lipid bilayers, thereby swelling the intercellular lipids^{130, 131}.

Conclusion

Skin permeation enhancement technology is a new and rapidly developing field which would significantly increase the number of candidates suitable for Transdermal Drug Delivery. Research in this area has proved the use of surfactant on the enhancement of permeation of drugs through skin. The techniques such as Differential scanning calorimeter, Fourier Transform Infra red, Nuclear magnetic resonance, Electron microscopy etc. have been very helpful in elucidating the mechanism of action and structure activity relationship of Penetration enhancers. Majority of studies reported indicate that the chemical structure of Penetration enhancers plays an important role on the permeation enhancement of drugs for some enhancers such as fatty acids, fatty alcohols and terpenes. Further studies are needed in the areas of evaluation of skin permeation enhancement vis-à-vis skin irritation in order to choose penetration enhancers which possess optimum enhancement effect with no skin irritation. A judicious selection of penetration enhancer would be very helpful in the successful development of topical and transdermal products.

Declaration

The author(s) declare that they have no competing interests or financial benefit from this work.

Acknowledgements

Authors are thankful to the principal KIET school of Pharmacy Ghaziabad.

References

1. Chein YW. Transdermal drug delivery & delivery systems. In: Yie W.Chein, ed. Novel drug delivery systems. New York: Marcel Dekker; 1992, 301-380.
2. Chein YW. Transdermal controlled systemic medications. New York: Marcel Dekker, INC. 1987.
3. Vyas SP, Khar RK. Transdermal drug delivery. In: Vyas SP & Khar RK, ed. Controlled drug delivery-concepts and advances. New Delhi, India: Vallabh Prakashan; 2002, 411-447.
4. Inpharmatechnologist.com – Rise of transdermal drug delivery technologies.htm- free newsletter.
5. Rein H. Experimental electroendosmotic studies on living human skin. *Z. Biol.* 81, 1924 124-128.
6. Champion RH, Burton JL, Burns DA & Breathnach SM. (eds). *Textbook of Dermatology* London: Blackwell Science; 1998.
7. Walker RB, Smith EW. The role of percutaneous penetration enhancers. *Adv Drug Deliv Rev* 1996; 18: 295-301.
8. Ayala-Bravo HA, Quintanar-Guerrero D, Naik A, Kalia YN, Cornejo-Bravo JM, Ganem Quintanar A. Effects of sucrose oleate and sucrose laurate on in vivo human stratum corneum permeability. *Pharm Res* 2003; 20: 1267–1273.
9. OECD Guidelines for the testing of chemicals, Section 4: Health Effects Test No. 428 Skin Absorption: In Vitro Method, Paris: OEC; 2004.
10. N.A. Monteiro-Riviere, Structure and function of skin, in: N.A. Monteiro-Riviere (Ed.), *Toxicology of the Skin: Target Organ Series*, Informa Healthcare USA, Inc., New York, 2010, pp. 1–18.
11. N.A. Monteiro-Riviere, Comparative anatomy, physiology and biochemistry of mammalian skin, in: D.W. Hobson (Ed.), *Dermal and Ocular Fundamentals and Methods*, Boca Raton: CRC Press; 1991, 3–71.
12. N.A.Monteiro Riviere, R.E.Baynes, J.E.Riviere, Animalskin morphology and dermal absorption, in: M.S. Roberts, K.A. Walters (Eds.), *Dermal Absorption and Toxicity Assessment*, second ed., USA: Informa Healthcare, Inc.; 2008, 17–35.
13. M.S. Roberts, A. Gierden, J.E. Riviere, N.A. Monteiro-Riviere, Solvent and vehicle effects on the skin, in: M.S. Roberts, K.A. Walters (Eds.), *Dermal Absorption and Toxicity Assessment*, second ed., Informa Healthcare USA, Inc., New York, 2007, pp. 433–447.
14. N.A. Monteiro-Riviere, Anatomical factors that affect barrier function, in: H. Zhai, K.P. Wilhelm, H.I. Maibach (Eds.), *Marzulli and Maibach's Dermatotoxicology*, seventh ed., CRC Press, New York, 2007, pp. 39–50.
15. P.M.Elias, Epidermallipids, barrier function, and desquamation, *J. Invest. Dermatol.* 80 (Suppl) (1983) 44s–49s.
16. M. Forster, M.A. Bolzinger, H. Fessi, S. Briancon, Topical delivery of cosmetics and drugs. Molecular aspects of percutaneous absorption and delivery, *Eur.J.Dermatol.* 19 (2009) 309–323.
17. Lavker, R.M. et al. (1986) Morphology of aged skin. In *Dermatological Clinics: The Aging Skin* (Lavker, R.M., ed.), pp. 1–20, Saunders, Philadelphia, PA, USA
18. Bouwstra JA, Dubbelaar FE, Gooris GS, Ponc M: The lipid organisation in the skin barrier. *Acta Derm Venereol Suppl* 2000; 208: 23–30.
19. Elias, P. M. & Feingold, K. R. Coordinate regulation of epidermal differentiation and barrier homeostasis. *Skin Pharmacol. Appl. Skin Physiol.* 14 (Suppl. 1), 28–34 (2001).
20. Menon GK. New insights into skin structure: scratching the surface. *Adv Drug Deliv Rev* 2002; 54: S3-17.
21. Williams AC. *Transdermal and topical drug delivery*. London: Pharmaceutical Press; 2003.
22. Kalbitz J, Neubert R, Wohlrab W: Modulation of drug penetration in the skin. *Pharmazie* 1996; 51: 619–637.

23. Ghosh,T.K. et al. (1997) Permeation enhancement. In *Transdermal and Topical Delivery Systems* (Ghosh,T.,Yum, S. and Pfister,W., eds), pp. 21–22, Interpharm Press, Buffalo Grove, IL, USA
24. Barry, B.W. (1991) Lipid-protein-partitioning of skin penetration enhancement. *J.Control.Release* 15, 237–248.
25. P. N. Gupta, V. Mishra, A. Rawat, P. Dubey, S. Mahor, S. Jain, D. P. Chatterji, and S. P. Vyas. Non-invasive vaccine delivery in transfersomes, niosomes and liposomes: A comparative study. *Int. J. Pharm.* 293(1–2):73–82 (2005).
26. Arellano, A., Santoyo, S., Martin, C., Ygartua, P., 1996.Enhancing effect of terpenes on the in vitro percutaneous absorption of diclofenac sodium. *Int. J. Pharm.* 130, 141-145
27. Maibach, H. et al. (1996) Epidermal enzymes as penetration enhancers in transdermal drug delivery. *J.Pharm.Sci.* 85, 249–262
28. Hofland, H.E. et al. (1994) Estradiol permeation from nonionic surfactant vesicles through human stratum corneum. *Pharm.Res.* 11, 659–664
29. Yokomizo,Y. (1996) Effect of phospholipids on the percutaneous penetration of drugs through the dorsal skin of the guinea pig in vitro. Part 3. Effects of phospholipids on several drugs having different polarities. *J.Control.Release* 42, 217–228
30. Masson, M. et al. (1999) Cyclodextrins as permeation enhancers: Some theoretical evaluations and in vitro testing. *Int.J.Pharm.* 59, 107–115.
31. Naito, S., Tominagaa, H., 1985. Percutaneous absorption of diclofenac sodium ointment. *Int. J. Pharm.* 24, 115?/124.
32. Yasukawa, T., Akiyoshi, Y., Hattori, A., Harada, S., 1985. Antiinflammatory topical formulations containing indomethacin and diclofenac. *Jpn. Kokai Tokkio Koho JP 60146823 A2*, August 22nd.
33. Nishihata, T., Kotera, K., Nakano, Y., Yamazaki, M., 1987. Rat percutaneous transport of diclofenac and influence of hydrogenated soya lecithin. *Chem. Pharm. Bull.* 35, 3807?/3812.
34. Williams, A.C., Barry, B.W., 1989. Essential oils as novel human skin penetration enhancers. *Int. J. Pharm.* 57, R7?/R9.
35. Takahashi, K., Tamagawa, S., Katagi, T., Yoshitomi, H., Kamada, A., Rytting, J., Nishihata, T., Mizuno, N., 1991a. In vitro transport of sodium diclofenac across rat abdominal skin: effect of selection of oleaginous component and the addition of alcohols to the vehicle. *Chem. Pharm. Bull.* 39, 154?/158.
36. Sarpotdar, R., Zatz, J.L., 1986a. Evaluation of penetration enhancement of lidocaine by nonionic surfactants through hairless mouse skin in vitro. *J. Pharm. Sci.* 75, 176?/181.
37. Sarpotdar, R., Zatz, J.L., 1986b. Percutaneous absorption enhancement by nonionic surfactants. *Drug Dev. Ind. Pharm.* 12, 1625?/1647.
38. Cappel, M.J., Kreuter, J., 1991. Effect of nonionic surfactants on transdermal drug delivery: I. Polysorbates. *Int. J. Pharm.* 69, 143?/153.
39. Iwasa, A., Irimoto, K., Kasai, S., Okuyama, H., Nagai, T., 1991. Effect of nonionic surfactants on percutaneous absorption of diclofenac sodium. *Yakuzaigaku* 51, 16?/21.
40. Lopez, A., Llinares, F., Cortell, C., Herraiez, M., 2000. Comparative enhancer effects of span 20 with Tween 20 and Azone on the in vitro percutaneous penetration of compounds with different lipophilicities. *Int. J. Pharm.* 202, 133-140.
41. Park, E., Chang, S.Y., Hahn, M., Chi, S., 2000. Enhancing effect of polyoxyethylene alkyl ethers on the skin permeation of ibuprofen. *Int. J. Pharm.* 209, 109?/119.
42. Shin, S.C., Cho, C.W., Oh, I.J., 2001. Effects of non-ionic surfactants as permeation enhancers towards piroxicam from the poloxamer gel through rat skins. *Int. J. Pharm.* 222, 199?/203.
43. Shokri, J., Nokhodchi, A., Dashbolaghi, A., Hassan-Zadeh, D., Ghafourian, T., Barzegar-Jalali, M., 2001. The effect of surfactants on the skin penetration of diazepam. *Int. J. Pharm.* 228, 99?/107.
44. Compendium of Chemical Terminology, 2nd ed. (the "Gold Book"). Compiled by A. D. McNaught and A. Wilkinson. Blackwell Scientific Publications, Oxford (1997). XML on-line corrected version: <http://goldbook.iupac.org> (2006-) created by M. Nic, J. Jirat, B. Kosata; updates compiled by A. Jenkins. ISBN 0-9678550-9-8. [doi:10.1351/goldbook](https://doi.org/10.1351/goldbook).
45. "Bubbles, Bubbles, Everywhere, But Not a Drop to Drink". *The Lipid Chronicles*.
46. D Attwood; AT Florence. *Surfactant Systems: Their Chemistry, Pharmacy and Biology*, Chapman and Hall Ltd., London, U. K., 1983, 21.
47. KA Walters. *Surfactants and percutaneous absorption*. In: *Prediction of Percutaneous Penetration: Methods, Measurements, Modelling*, Scott, R. C., Guy R. H., Hadgraft, J. (Eds). IBC Technical Services Ltd., London, U. K., 1990, 148-162.
48. R.J. Scheuplein, L. Ross, Effects of surfactants and solvents on the permeability of epidermis, *Journal of the Society of Cosmetic Chemists*. 1970;21(13): 853-873.
49. Froebe C. L., Simion F. A., Rhein L. D., Cagan R. H., Kligman A.(1990) Stratum corneum lipid removal by surfactants: relation to in vivo irritation. *Dermatologica* 181:277–283.
50. Leveque J. L., de Rigal J., Saint-Leger D., Billy B.(1993) How does sodium lauryl sulfate alter the skin barrier function in man? A multiparametric approach. *Skin Pharmacol.* 6:111–115.
51. Patil S., Singh P., Maibach H.(1995) Radial spread of sodium lauryl sulfate after topical application. *Pharm. Res.* 12:2018–2023.
52. Patil S., Singh P., Sarasour K.Maibach H.(1995) Quantification of sodium lauryl sulfate penetration into the skin and underlying tissue after topical application, pharmacological and toxicological implications. *J. Pharm. Sci.* 84:1240–1244.
53. Rhein, L.D., Robbins, C.R., Fernee, K., Cantore, R., 1986. Surfactanta structure effects on swelling of

- isolated human stratum corneum. *J. Soc. Cosmet. Chem.* 37, 199?/210.
54. Gibson, K.T., Teall, M.R., 1983. Interactions of C12 surfactants with the skin: changes in enzymes and visible and histological features of rat skin treated with sodium lauryl sulphate. *Food Chem. Toxicol.* 21, 587?/594.
 55. Kitagawa S., Kasamaki M., Ikarashi A., Effect of n-Alkyltrimethylammonium on skin permeation of Benzoic acid through excised Guinea pig dorsal skin. G0504A ISSN:0009-2363 *Chem. Pharm. Bull.*, 48, 1698—1701 (2000).
 56. Breuer, M.M., 1979. The interaction between surfactants and keratinous tissues. *J. Soc. Cosmet.* 30, 41?/64.
 57. Walters, K.A., Walker, M. and Olejnik, O. Non-ionic surfactant effects on hairless mouse skin permeability characteristics. *J. Pharm. Pharmacol.* 40,525-9 (1988).
 58. N. Kanikkannan, K. Kandimalla, S.S. Lamba, M. Singh, Structure–activity relationship of chemical penetration enhancers in transdermal drug delivery, *Curr. Med. Chem.* 7 (6) (2000) 593–608.
 59. H. Tanojo, E. Boelsma, H.E. Junginger, M. Ponec, H.E. Bodde, In vivo human skin barrier modulation by topical application of fatty acids, *Skin Pharmacol. Appl. Skin Physiol.* 11 (2) (1998) 87–97.
 60. B.J. Aungst, Structure/effect studies of fatty acid isomers as skin penetration enhancers and skin irritants, *Pharm. Res.* 6 (1989) 244–247.
 61. Aungst, B. J., Rogers, N. J. and Shefter, E. 1986. Enhancement of naloxone penetration through human skin in vitro using fatty acids, fatty alcohols, surfactants, sulfoxides and amines. *International Journal of Pharmaceutics.* 33, 225-234.
 62. S. Mitragotri, Synergistic effect of enhancers for transdermal drug delivery, *Pharm. Res.* 17 (2000) 1354–1359.
 63. E.H. Choi, S.H. Lee, S.K. Ahn, S.M. Hwang, The pretreatment effect of chemical skin penetration enhancers in transdermal drug delivery using iontophoresis, *Skin Pharmacol. Appl. Skin Physiol.* 12 (1999) 326–335.
 64. S. Chesnoy, D. Durand, J. Doucet, G. Couarraze, Structural parameters involved in the permeation of propranolol HCl by iontophoresis and enhancers, *J. Control. Release* 58 (1999) 163–175.
 65. L. Wearley, Y.W. Chien, Enhancement of the in vitro skin permeability of azidothymidine (AZT) via iontophoresis and chemical enhancer, *Pharm. Res.* 7 (1990) 34–40.
 66. Heather A.E. Benson. Transdermal drug delivery: penetration enhancement techniques. *Curr. Drug Delivery.* 2005;2:23-33.
 67. Laughlin, R.G., 1978. Relative hydrophilicities among surfactants hydrophilic groups. In: Brown, G.H. (Ed.), *Advances in Liquid Crystals.* Academic Press, New York.
 68. Israelachvili, J. N. and Wennerström, H., *J. Phys. Chem.*, 1992. 96, 520.
 69. Chowhan, Z. T. and Pritchard, R. 1978. Effect of surfactants on the percutaneous absorption of naproxen. I. Comparison of rabbit, rat and human excised skin. *Journal of Pharmaceutical Sciences.* 67, 1272-1274.
 70. Nokhodchi, A., Shokri, J., Dashbolaghi, A., Hassan-Zadeh, D., Ghafourian, T. and Barzegar-Jalali, M. 2003. The enhancement effect of surfactants on the penetration of lorazepam through rat skin. *International Journal of Pharmaceutics.* 250, 359-369.
 71. Kushla, G. P., Zatz, J. L., Mills, O. H. J. and Berger, R. S. 1993. Noninvasive assessment of anaesthetic activity of topical lidocaine formulations. *Journal of Pharmaceutical Sciences.* 82, 1118-1122.
 72. P.C. Rigg, B.W. Barry, Shed snake skin and hairless mouse skin as model membranes for human skin during permeation studies, *J. Invest. Dermatol.* 94 (1990) 235–240.
 73. Augiar, A.J., Weiner, M.A., 1969. Percutaneous absorption of chloramphenicol solution. *J. Pharm. Sci.* 58, 210?/215.
 74. Ribaud, C.H., Garson, J.C., Doucet, J., Leveque, J.L., 1994. Organisation of stratum corneum lipids in relation to permeability: influence of sodium lauryl sulphate and preheating. *Pharm. Res.* 11, 1414?/1418.
 75. Goodman, M., Barry, B., 1989. Action of penetration enhancers on human stratum corneum as assessed by differential scanning calorimetry. *Transdermal Drug Deliv.* 35, 574?/593.
 76. Borrás-Blasco, J., Lopez, A., Morant, M.J., Diez-Sales, O., Herraiz-Domingues, M., 1997. Influence of sodium lauryl sulphate on the in vitro percutaneous absorption of compounds with different lipophilicity. *Eur. J. Pharm. Sci.* 5, 15-22.
 77. Darshan K. Parikh, Tapash K. Ghosh, Feasibility of Transdermal Delivery of Fluoxetine. *AAPS PharmSciTech* 2005; 6 (2), 22
 78. K. Bhaskar Reddy, M.Jyostna, N.Audinarayana, C.Madhavi Latha, E.Mohanambal. The Enhancement Effect of Surfactants on the Penetration of Nitrendipine through Rat Skin. *Indian Journal of Novel Drug delivery* 4(1), Jan-Mar, 2012, 38-42.
 79. Sang-Chul Shin, Cheong-Weon Cho, In-Joon Oh Effects of non-ionic surfactants as permeation enhancers towards piroxicam from the poloxamer gel through rat skins. *International Journal of Pharmaceutics* 222 (2001) 199–203
 80. Jocelyne Piret, Andre´De´Sormeaux, He´Le`Ne Cormier, Julie Lamontagne, Pierrette Gourde, Julianna Juha´Sz, Michel G. Bergeron. Sodium Lauryl Sulfate Increases the Efficacy of a Topical Formulation of Foscarnet against Herpes Simplex Virus Type 1 Cutaneous Lesions in Mice. *Antimicrobial Agents And Chemotherapy*, Sept. 2000, p. 2263–2270.
 81. Sankar V, Praveen C, Prasanth KG, Srinivas CR, Ruckmann K. Formulation and evaluation of a proniosome hydrocortisone gel in comparison with a commercial cream. *Pharmazie* 009; 64:731–4.
 82. C.-W. Cho, J.-S. Choi, and S.-C. Shin, “Enhanced local anesthetic action of mepivacaine from the bioadhesive gels,” *Pakistan Journal of*

- Pharmaceutical Sciences, vol. 24, no. 1, pp. 87–93, 2011.
83. Ekapol Limpongsa, Kraisri Umprayn, Preparation and Evaluation of Diltiazem hydrochloride Diffusion-Controlled Transdermal Delivery System. *AAPS PharmSciTech*, Vol. 9, No. 2, June 2008.
 84. Adamo Fini, Valentina Bergamante, Gian Carlo Ceschel, Celestino Ronchi, and Carlos Alberto Fonseca De Moraes, Control of Transdermal Permeation of Hydrocortisone Acetate from Hydrophilic and Lipophilic Formulations. *AAPS PharmSciTech*, Vol. 9, No. 3, September 2008.
 85. Ganem-Quintanar, Lafforgue AC, Falson-Rieg F, Buri P. Evaluation of the transepidermal permeation of diethyleneglycol monoethylether and skin water loss. *Int J Pharm*. 1997;147: 165–72.
 86. Godwin A, Kim NH, Felton LA. Influence of Transcutol CG on the skin accumulation and transepidermal permeation of ultraviolet absorbers. *Eur J Pharm Biopharm*. 2002;53:23–7.
 87. Shishu, Sunita Rajan, Kamalpreet, Development of Novel Microemulsion-Based Topical Formulations of Acyclovir for the Treatment of Cutaneous Herpetic Infections, *AAPS PharmSciTech*, Vol. 10, No. 2, June 2009 DOI: 10.1208/s12249-009-9242-1
 88. Constantinides PP, Scalart JP, Lancaster C, Marcello J, Marks G, Ellens H, Smith PL. Formulation and intestinal absorption enhancement evaluation of water-in-oil microemulsions incorporating medium-chain glycerides. *Pharm Res* 1994;11:1385–90.
 89. Lopes LB, Murphy N, Nornoo A. Enhancement of transdermal delivery of progesterone using medium-chain mono and diglycerides as skin penetration enhancers. *PharmDevTechol*. 2009. Doi :10.1080/10837450902814180.
 90. Lopes LB, Collett JH, Bentley MV. Topical delivery of cyclosporin A: an in vitro study using monoolein as a penetration enhancer. *Eur J Pharm Biopharm* 2005;60:25–30.
 91. Herai H, Gratieri T, Thomazine JA, Bentley MV, Lopez RF. Doxorubicin skin penetration from monoolein-containing propylene glycol formulations. *Int J Pharm* 2007;329:88–93.
 92. Magnusson BM, Cross SE, Winckle G, Roberts MS. Percutaneous absorption of steroids: determination of in vitro permeability and tissue reservoir characteristics in human skin layers. *Skin Pharmacol Physiol* 2006;19:336–42.
 93. Frum Y, Khan GM, Sefcik J, Rouse J, Eccleston GM, Meidan VM. Towards a correlation between drug properties and in vitro transdermal flux variability. *Int J Pharm* 2007;336:140–7.
 94. Zar T, Graeber C, Perazella MA. Recognition, treatment, and prevention of propylene glycol toxicity. *Semin Dial* 2007;20:217–9.
 95. Mrunali R. Patel, Rashmin B. Patel, Jolly R. Parikh, Ajay B. Solanki, and Bharat G. Patel Effect of Formulation Components on the In Vitro Permeation of Microemulsion Drug Delivery System of Fluconazole, *AAPS PharmSciTech*, Vol. 10, No. 3, September 2009 DOI:10.1208/s12249-009-9286-2
 96. Obata Y, Takayama K, Machida Y, Nagai T. Combined effect of cyclic monoterpenes and ethanol on percutaneous absorption of diclofenac sodium. *Drug Des Discov*. 1991;8:137–44.
 97. Adnan Azeem,1,2Farhan J. Ahmad,1Roop K. Khar,1and Sushama Talegaonkar, Nanocarrier for the Transdermal Delivery of an Antiparkinsonian Drug *AAPS PharmSciTech*, Vol. 10, No. 4, December 2009 DOI: 10.1208/s12249-009-9306-2
 98. Shah VP. Skin penetration enhancers: scientific perspectives. In: Hsieh DS, editor. *Drug permeation enhancement; theory and applications*. New York: Marcel Dekker; 1994. p. 19–24.
 99. Mahmoud Mokhtar Ahmed Ibrahim, Omaima A. Sammour, Mohamed A. Hammad, and Nagia A. Megrab. In Vitro Evaluation of Proniosomes as a Drug Carrier for Flurbiprofen. *AAPS PharmSciTech*, Vol. 9, No. 3, September 2008 DOI: 10.1208/s12249-008-9114-0.
 100. Aliaa Nabil ElMeshad and Mina Ibrahim Tadros Transdermal Delivery of an Anti-Cancer Drug via W/O Emulsions Based on Alkyl Polyglycosides and Lecithin: Design, Characterization, and In Vivo Evaluation of the Possible Irritation Potential in Rats. *AAPS PharmSciTech*, Vol. 12, No. 1, March 2011 DOI: 10.1208/s12249-010-9557-y
 101. Friedman-Kien, A. E., R. J. Klein, R. D. Glaser, and S. M. Czelusniak. 1986. Treatment of recurrent genital herpes with topical alpha interferon combined with nonoxyl 9. *J. Am. Acad. Dermatol*. 5:989-994.
 102. Endo M, Yamamoto T, Ijuin T. Effect of nonionic surfactants on the percutaneous absorption of tenoxicam. *Chem Pharm Bull*. 1996;44: 865–867.
 103. Okuyama H, Ikeda Y, Kasai S, Imamori K, Takayama K, Nagai T. Influence of non-ionic surfactants, pH and propylene glycol on percutaneous absorption of piroxicam from cataplasm. *Int J Pharm*. 1999;186: 141–148.
 104. Shufang Nie, WL Wendy Hsiao, Weisan Pan, Zhijun Yang, Thermoreversible Pluronic® F127-based hydrogel containing liposomes for the controlled delivery of paclitaxel: in vitro drug release, cell cytotoxicity, and uptake studies. *International Journal of Nanomedicine* 2011(6):151-166.
 105. Ghassan Z Abdullah et al, In vitro permeation and in vivo anti-inflammatory and analgesic properties of nanoscaled emulsions containing ibuprofen for topical delivery. *International Journal of Nanomedicine*. 2011;6 :387–396.
 106. Ji-hui Zhao, Li Ji, hui Wang, Zhi-Qiang chen, Yong-Tai Zhang, Ying Liu, Nian-Ping Feng. Microemulsion-based novel transdermal delivery system of tetramethylpyrazine: preparation and evaluation in vitro and in vivo. *International Journal of Nanomedicine* 2011;6 1611–1619.
 107. Ibrahim A. Alsarra, A.A. Bosela, S.M. Ahmed, G.M. Mahrous, Proniosomes as a drug carrier for transdermal delivery of ketorolac. *European*

- Journal of Pharmaceutics and Biopharmaceutics 59 (2005) 485–490.
108. Ancel HC, Allen LV, Popovich NG. Pharmaceutical dosage forms and drug delivery systems. 5th ed., New Delhi: Lippincott Williams & Wilkins; 2005, p. 311-312.
109. Hemangi J Patel *1, Jitendra S Patel 2, B G Desai 3, Keyur D Patel, Permeability studies of anti hypertensive drug amlodipine besilate for transdermal delivery. Asian Journal of Pharmaceutical and Clinical Research. Vol.3 Issue 1, January-March 2010, 31-34.
110. Gannu RP, Chinna R. Yamsani V, Yamsani SK, Yamsani MR. Enhanced bioavailability of Lacidipine via microemulsion based transdermal gel: formulation optimization, ex- vivo and in- vivo characterization. International Journal of pharmaceutics, 2010; 388;1(2):231-241.
111. Pao-Chu W, Yei-Huan L, Jui Sheng C, Yaw-Bin H, Yi-Hung T. The effect of component of microemulsion for transdermal delivery of nicardipine hydrochloride. Drug Development and Industrial Pharmacy, 2010; 36; 12: 1398-1403.
112. Sheikh SN, Faiyaz S, Sushma T, Javed A. Formulation Development and Optimization using Nanoemulsion Technique: A Technical Note. AAPS Pharmaceutical Science and Technology, 2007; 8(2): E1-E6.
113. Inayat Bashir Pathan, C.Mallikarjuna Setty, Enhancement of Transdermal Delivery of Tamoxifen Citrate using Nanoemulsion Vehicle. International Journal of PharmTech Research CODEN (USA): IJPRIF ISSN : 0974-4304 Vol. 3, No.1, pp 287-297, Jan-Mar 2011
114. Mukherjee B, Kanupriya, Mahapatra S, Das S, Patra B: Sorbitan monolaurate 20 as a potential skin permeation enhancer in transdermal patches. J Appl Res 2005; 1: 96–108.
115. Shin SC, Choi JS: Enhanced efficiency of triprolidine by transdermal application of the EVA matrix system in rabbits and rats. Eur J Pharm Biopharm 2005; 61: 14–19.
116. T. Ogiso, M. Shintani, Mechanism for the enhancement effect of fatty acids on the percutaneous absorption of propranolol, J. Pharm. Sci. 79 (12) (1990) 1065–1071.
117. B.M. Elyan, M.B. Sidhom, F.M. Plakogiannis, Evaluation of the effect of different fatty acids on the percutaneous absorption of metaproterenol sulfate, J. Pharm. Sci. 85 (1) (1996) 101–105.
118. Kenneth A. Walters, W. Bialik, Keith R. Brain, The effects of surfactants on penetration across the skin. International Journal of Cosmetic Science 15,260-270 (1993).
119. M.J. Pikal, The role of electroosmotic flow in the transdermal iontophoresis, Adv. Drug Deliv. Rev. 46 (1–3) (2001) 281–305.
120. H.E.J. Hofland, J.A. Bouwstra, F. Spies, H.E. Bodde, J.F. Nagelkerke, C. Cullander, H.E. Junginger, Interactions between non-ionic surfactant vesicles and human stratum corneum in vitro: freeze fracture scanning microscopy and confocal laser scanning microscopy, J. Liposome Res. 5 (1995) 241–263.
121. Karande, P., Jain, A. and Mitragotri, S. 2004. Discovery of transdermal penetration enhancers by high-throughput screening. Nature Biotechnology. 22, 192-197.
122. Chowhan, Z. T. and Pritchard, R. 1978. Effect of surfactants on the percutaneous absorption of naproxen. I. Comparison of rabbit, rat and human excised skin. Journal of Pharmaceutical Sciences. 67, 1272-1274.
123. Vaddi, H.K., Wang, L.Z., Ho, P.C., Chan, S.Y., 2001. Effects of some enhancers on the permeation of haloperidol through rat skin in vitro. Int. J. Pharm. 212, 247-255.
124. Megwa S. A., Cross S. E., Benson H. A. E., Roberts M. S., J. Pharm. Pharmacol., 52, 919–928 (2000).
125. Kitagawa S., Kasamaki M., Ikarashi A., Chem. Pharm. Bull., 48,1698–1701 (2000).
126. Smith J. C., Irwin W. J., Int. J. Pharmaceut., 210, 69–82 (2000).
127. James-Smith, M. A., D. Shekhawat, and D. O. Shah. Importance of micellar lifetime and sub-micellar aggregates in detergency processes. Tenside Surf. Deterg. 44(3):142–154, 2007.
128. Patist, A., J. R. Kanicky, P. K. Shukla, and D. O. Shah. Importance of micellar kinetics in relation to technological processes. J. Colloid Interface Sci. 245(1):1–15, 2002.
129. U.F. Schaefer, K. Elbert, F. Ruchatz Testing Of Different Substances For Their Dermal Penetration Enhancement. Department of Biopharmaceutics and Pharm. Technology, Saarland University, Im Stadtwald, D-66123 Saarbrücken, GERMANY
130. Ganem-Quintanar, Lafforgue AC, Falson-Rieg F, Buri P. Evaluation of the transepidermal permeation of diethyleneglycol monoethylether and skin water loss. Int J Pharm. 1997;147:165–72.
131. Godwin A, Kim NH, Felton LA. Influence of Transcutol CG on the skin accumulation and transepidermal permeation of ultraviolet absorbers. Eur J Pharm Biopharm. 2002;53:23–7.